

Doxem[®] Precise Dry Flowable Cockroach Bait

Service Protocol 3: Managing German Cockroaches Indoors in Schools

INITIAL TREATMENT: ONCE ANNUALLY

Consult with the IPM coordinator/administration to determine where pest populations have been noted and where application of pesticides is allowed. Be sure to determine Doxem Precise is included on their approved products list and **you follow** any local requirements for applying bait products in schools. Be sure to follow any notification protocols required in your service area.

Complete a thorough inspection of the property and properly identifying the pest in question. Begin by assessing the population size, location and severity. Note conducive conditions and recommend repair/mitigation of these conditions where appropriate.

Be sure to make applications of Doxem ONLY in unoccupied areas. Application in classrooms is only allowed when occupants are NOT present.

Areas other than classrooms:

STEP 1: Physically remove as many cockroaches as possible using integrated means such as vacuuming, glue boards or traps.

STEP 2: In the kitchen, only make applications to cracks/crevices. Be sure to avoid contaminating and food/feed products or preparation surfaces, dishes or utensils at the time of treatment. Make applications of 3-8 seconds where cockroaches are observed. Examples of areas to treat:

- Cabinet voids
- High moisture areas (dishwashers, sinks)
- Refrigerators (motor housings, compressors, drains)
- Around ovens/microwaves
- Cardboard box storage

Doxem[®] Precise Dry Flowable Cockroach Bait

In classrooms:

STEP 1: Be sure the room is unoccupied before treatment.

STEP 2: Physically remove as many cockroaches as possible using integrated means such as vacuuming, glue boards or traps.

STEP 3: Make applications of Doxem Precise. Apply 3-8 seconds into voids, cracks & crevices where cockroaches are observed. Examples of areas to treat:

- Cabinet voids
- Bathroom/sink areas
- Storage closets/voids
- Cubbies/coat closets
- Other areas where cockroaches are observed

FOLLOW UP TREATMENTS:

STEP 1: Re-inspect the property and all treated areas. Determine current activity level and areas where cockroaches might still be active. If the area was missed on the initial treatment, apply Doxem Precise to those areas at the 3-8 seconds per void/crack/crevice. In other areas, apply Doxem only in areas where cockroaches likely consumed some/all of the bait material.

QUARTERLY & AS NEEDED TREATMENTS:

STEP 1: Re-inspect the property and all treated areas. Determine current activity level and areas where new infestations of cockroaches might be active. If the area was not treated during the initial treatment, apply Doxem Precise using the PDS to those areas at the 3-8 seconds per void/crack/crevice. In other areas, apply Doxem only in areas where cockroaches likely consumed some/all of the bait material.

For More Information:

CSI-PDS.com



CSI-Pest.com
f @csipest t @csi_pest