



# QUALI-PRO CONTRADO® SC

EPA REG. NO. 53883-534

EXPIRES: DEC 31, 2028

**Control Solutions Inc.**  
**5903 Genoa Red Bluff**  
**Pasadena, Texas, 77507**  
**(800) 242-5562**

## ***PRODUCT BULLETIN***

**STATES:** AK, AL, AR, AZ, CO, CT, DC, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NV, OH, OK, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label.

**This product bulletin must be in the possession of the user at the time of application of the product.** All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### **ADDITIONAL PESTS OF ORNAMENTAL PLANTS IN OUTDOOR LANDSCAPES, INTERIOR PLANTSCAPES, AND COMMERCIAL GREENHOUSES AND NURSERIES**

<b>Ornamentals in outdoor landscapes and interior plantscapes and commercially produced in indoor nurseries, outdoor nurseries (both field-grown and containerized plants), greenhouses, shade houses, lath houses, and hoop houses.</b>		
Bulb, corm and tuber crops (such as tulips, calla lilies) Evergreens, including conifers Flowering plants Foliage plants	Ground covers Non-bearing fruit and nut trees <sup>1</sup> Ornamental grasses Ornamental trees Palms	Perennial plants Shrubs Succulents Vines (non-bearing) <sup>1</sup>
<sup>1</sup> Non-bearing fruit and nut trees and vines are those trees and vines that will not bear edible fruit or nuts for one year after application.		

Contrado Ornamental Foliar Application		
Target Pest	Contrado Dilution Rate per 100 gallons	lb. ai per 100 gallons
Fall Webworm, Spongy Moth Caterpillar ( <i>Formerly Gypsy Moth</i> )	1 – 8 fl. oz.	0.013 – 0.104
Sawfly Larvae	2 – 4 fl. oz.	0.026 – 0.052
European Pepper Moth Japanese Beetle Adults	4 – 10 fl. oz.	0.052 – 0.13
Viburnum Leaf Beetle	4 – 16 fl. oz.	0.052 – 0.208
Zimmerman Pine Moth	16 fl. oz.	0.208
Banded Ash Clearwing Borer Moth <sup>2</sup> ( <i>Podosesia aureocincta</i> ) <sup>2</sup>	32 fl. oz.	0.416
<sup>2</sup> Trunk Spray Application <ul style="list-style-type: none"> <li>• Use sufficient water volume to provide good coverage of target foliage without excessive runoff.</li> <li>• Use the lower labeled rates when targeting smaller or younger and more vulnerable pest populations.</li> <li>• Apply higher rates when pest populations are more mature or under heavier pressure to ensure effective control.</li> </ul>		

Contrado Ornamental Soil Drench Application		
Target Pest	Contrado Dilution Rate per inch DBH of trees or foot of height of shrubs	lb. ai per inch DBH of trees or foot of height of shrubs
Flatheaded Apple Tree Borer	0.25 fl. oz.	0.0032
Peachtree Borer		
Rhododendron Lace Bug		
Hemlock Woolly Adelgid		
False Oleander Scale		
Magnolia Scale		
Oriental Beetle Larvae		
<ul style="list-style-type: none"><li>• For optimal control, apply the product to the soil 45–60 days before anticipated pest hatch or emergence to allow adequate systemic uptake by the plant. In most regions, applications made in April to May are effective; however, users should consult local Extension services for region-specific timing.</li><li>• Apply within 1–3 feet of the root flare of trees and shrubs, using enough volume to reach a majority of the root zone and promote systemic absorption.</li></ul>		